Mass-gathering Health Research Foundational Theory: Part 2 - Event Modeling for Mass Gatherings

Sheila A. Turris, NP, PhD; Adam Lund, MD, MEd, FRCPC; Alison Hutton, RN, PhD; Ron Bowles, EMA II,

PhD;³ Elizabeth Ellerson, MPH;⁴ Malinda Steenkamp, PhD; Jamie Ranse, RN, FRCNA, FACN;⁵

Paul Arbon, RN, PhD

- 1. Department of Emergency Medicine, University of British Columbia, Vancouver, British Columbia, Canada
- 2. School of Nursing, University of Victoria, Victoria, British Columbia, Canada
- 3. Justice Institute of British Columbia, New Westminster, British Columbia, Canada
- 4. Flinders University, World Health Organization Collaborating Centre for Mass Gatherings and High Consequence/High Visibility Events, Flinders University of South Australia, Adelaide, South Australia, Australia
- 5. University of Canberra, Faculty of Health, Bruce, Australian Capital Territory, Australia

Abstract

Background: Current knowledge about mass gathering health (MGH) fails to adequately inform the understanding of mass gatherings (MGs) because of a relative lack of theory development and adequate conceptual analysis. This report describes the development of a series of event lenses that serve as a beginning "MG event model," complimenting the "MG population model" reported elsewhere.

Methods: Existing descriptions of "MGs" were considered. Analyzing gaps in current knowledge, the authors sought to delineate the population of events being reported. Employing a consensus approach, the authors strove to capture the diversity, range, and scope of MG events, identifying common variables that might assist researchers in determining when events are similar and might be compared. Through face-to-face group meetings, structured breakout sessions, asynchronous collaboration, and virtual international meetings, a conceptual approach to classifying and describing events evolved in an iterative fashion.

Findings: Embedded within existing literature are a variety of approaches to event classification and description. Arising from these approaches, the authors discuss the interplay between event demographics, event dynamics, and event design. Specifically, the report details current understandings about event types, geography, scale, temporality, crowd dynamics, medical support, protective factors, and special hazards. A series of tables are presented to model the different analytic lenses that might be employed in understanding the context of MG events.

Interpretation: The development of an event model addresses a gap in the current body of knowledge *vis* a *vis* understanding and reporting the full scope of the health effects related to MGs. Consistent use of a consensus-based event model will support more rigorous data collection. This in turn will support meta-analysis, create a foundation for risk assessment, allow for the pooling of data for illness and injury prediction, and support methodology for evaluating health promotion, harm reduction, and clinical response interventions at MGs.

Introduction

Less than 25 years old, mass-gathering health (MGH) is a relatively new field of research.¹ As such, the science underpinning this body of knowledge is young and developing rapidly.² Current knowledge, however, fails to adequately inform the understanding of mass gatherings (MGs) because of the lack of theory development and adequate conceptual analysis.³ In December of 2013, the World Health Organization (WHO; Geneva, Switzerland) Collaborating Center for Mass Gatherings and High Risk/Visibility Events at Flinders University, South Australia, hosted a scientific meeting bringing together international MGH researchers. Representatives were from: Flinders

Scenario 1

You are the medical director for a four-day music festival held in a rural setting, annually, in summer. There are 1,200 people living in the community. The forecast is for hot, dry, windy conditions. The event will attract more than 30,000 daily attendees, many of whom will camp overnight. The nearest hospital is a 2-hour drive away.

Scenario 2

You are a physician in an urban emergency department. A qualifying marathon is going to be held in your city on Sunday. The forecast is for rain and cool temperatures. The event will attract more than 40,000 participants and an unknown number of spectators. There are three hospitals within a 20-minute drive. A bomb exploded at a similar event two months earlier, which has media, police, and health officials on "high alert."

Table 1. Event Scenarios

Turris & 2014 Prehospital and Disaster Medicine

University and University of Canberra, in Australia; the University of British Columbia (BC) and the Justice Institute of BC, in Canada; the WHO; and Public Health England (London, England, United Kingdom).

The MGH Collaborating Team, formed during this meeting, discussed various models related to MG populations and events to support a common understanding about their human health effects. The goal of a larger, international consensus project is the creation of a minimum data set and data dictionary for MGH research. This will support description, measurement, comparison, evaluation, and reporting on parameters of interest, permitting international comparisons that, to date, has been impossible.^{2,4-6}

Mass-gathering events are complex phenomena involving the interaction between a number of elements. This manuscript is the second of two related reports that explore key elements of MGH research. The first paper⁷ concerns the various *populations of people* involved in MGs and proposes a MGH population model. The current manuscript complements this work by describing and characterizing *MG events,* identifying specific variables of interest to MG researchers, and addressing a gap in the current body of knowledge with regard to understanding the full scope of MGs and their health effects.

A robust event model is increasingly important as some events attract international participants and attendees in large numbers. In the interactions between members of the host and event populations, there exist opportunities for injuries,^{8,9} the spread of infectious disease (ID),¹⁰⁻¹⁷ disorderly crowd behavior and movement,¹⁷⁻²¹ and the overflow of ill and injured from the event to the surrounding community during a mass-casualty incident, perhaps overwhelming the host infrastructure.^{20,22-26}

Consistent data collection with regard to events will potentially:(1) permit meta-analysis and pooling of data sources internationally;(2) further develop the methodology for evaluating health promotion and harm reduction efforts; (3) create a solid foundation for risk assessment and illness and injury prediction modeling; and (4) provide operational support for front-line clinicians with regard to planning for on-site medical teams.^{3,5,6,9}

Methods

The authors critically appraised existing MG definitions and descriptions in published literature. When it became apparent that published definitions lacked sufficient breadth and depth to support the group in defining, describing, and explaining the health effects of MGs, a series of tables with event characteristics were developed in an iterative fashion, capturing both academically- and operationally-relevant lenses for MGs.

Findings and Interpretation

Characterizing MG Events

An ongoing challenge for MG researchers and organizers is how to categorize events in a way that allows for grouping, comparing, and contrasting findings. The MG literature covers a staggering range of events, each with a number of event-related characteristics.

The cases above (Table 1) are similar in that large groups (ie, 30-40,000 people) will gather for a specific event, but their divergent characteristics are far more intriguing (ie, one is a 1-day sporting event, the other a multi-day cultural event). One is in a small community with limited local health care capacity, while the other is a large community with a substantial pre-existing medical system.

How might these differences affect illness and injury rates? Can injury and illness rates be reduced? How will treatment planning differ for individuals who present for care? A careful read of existing event descriptions²⁷⁻⁵⁵ reveals a multitude

of event-related variables that are inconsistently reported and currently lack clear definitions (Table 2). The results of the team's iterative process are presented below. Macro-level characteristics explore the interplay between the place, people, and the event itself, informing methodology, data collection, and meta-analysis. Meso-level characteristics include the "demographics" (ie, the immutable parts of the event), the "dynamics" (ie, what people do; how they feel and react), and the "design" (ie, what event planners/ producers do to protect attendees/participants from illness and injury) of MGs and their relationship to risk assessment and predictive modeling. Micro-level characteristics inform planning and mitigation efforts of frontline clinicians, medical teams, and other event personnel. Micro-level characteristics will be detailed in a future publication regarding the consensus-based minimum data set and data dictionary.

Below, the authors present a broader explanation of MG demographics, dynamics, and design. The elements in this discussion represent various variables of interest to MG researchers that may be useful for risk assessment and predictive modeling.

Demographics of an Event

In existing literature, common starting points for grouping and characterizing MG events include event type, geography, size, and temporal considerations.

Event Type—With the exception of relatively-common events, such as marathons or cycling events, event types are typically

Event Dimension	Category	Classification Examples	Citations
Demographics	Event Type	Races, cycling events, religious pilgrimages, cultural events, and music festivals.	,18, 26-34
	Geography	Terrain, bounded or unbounded, shifting or fixed footprint, site access and egress, multi or single venue.	17, 26, 28-30, 35
	Size	Number of attendees/participants, multi-event versus single.	26, 36
	Temporality	Duration (days, hours), time of year, time of day, season, recurrent annual event vs first-time event, peak time of attendance.	26, 28-29, 34-35
Dynamics	Crowd Type	Gender mix, age mix, families, disabilities, special populations.	26, 28, 29
	Crowd Behavior	Mood, density, activity levels, queuing, movement, behavior, predispositions, motivations, crowd movement, and flow.	17, 33, 36-42
	Purpose of Event	Sport, arts, religious, political, mixed; protests, riots, celebrations.	20, 26-27, 29
	Political Context	Within host community/country, police state versus democracy, civic versus private, profit versus not for profit.	26-27, 29
Design	Protective Factors	Crowd resilience, health promotion, illness prevention, police/ security onsite.	24, 28-29, 43-46
	Special Hazards	Climate, weather conditions, motorization, obstacle course, infectious disease exposure, alcohol and drug use, mosh pits, fireworks.	19, 20, 26, 29, 31-36, 39, 47- 49
	Onsite Health Services	First aid only, higher level of care, location, deployment, signage, triage.	24, 26, 28, 31, 35, 39, 41, 50-52
	Host Community Burden	Mitigation of event related increases in volume for host community infrastructure.	24, 43, 53-54

 Table 2. Dimensions of an Event Model as Reported in the Literature

Turris & 2014 Prehospital and Disaster Medicine

recorded without reference to a category. That is, every event is treated as unique. However, Turris et al argued that

an initial typology of events would include sporting, arts-related, religious/political, and miscellaneous events.²¹ This is a macro-level model, which has limited utility for developing nuanced explanatory and predictive models. For example, it does not distinguish between contact and noncontact sporting events, spectator vs participant sporting events, and so on. It does, however, present an initial "high-level" categorization as a starting point for grouping and comparing various types of MGs.

Strictly speaking, the event type (eg, fireworks display, boat show, and Papal visit) is only one piece of data that will help researchers and clinicians determine the degree to which two different events may be alike. Other parameters, such as whether or not an event is primarily spectator- or participant-based, may be as helpful in selecting events for comparison.

Geography—Every event has a geographical character and site-specific descriptors. For example, at a local music festival held annually in Vancouver, BC (Canada), the grounds include a lake, raising the specter of accidental drowning. Similarly,

at the Suwa Onbashira Festival held every six years in Japan, men ride logs down steep slopes.³⁴ The geography of both events described creates a set of safety issues for event planners, attendees, and on-site medical services.

Bounded, Unbounded, or Compound Events—Milsten et al identified that an event with a mobile population would have a different risk profile than an event for which spectators were seated throughout the event.⁵⁰ This is, in part, due to the fact that in unbounded events, there is no control with regard to the capacity of the venue. In contrast, for an event held at a stadium with fixed seats, event planners have some control over the number of participants and spectators. A variety of event "footprints" that might be of interest to researchers and clinicians are presented in Table 3.

Considerations with regard to event boundaries are reported in the literature. For example, marathons tend to have a shifting footprint. As runners continue along the course, the early parts of the course are closed. This has implications for medical

deployment as health care may be offered by both fixed and mobile response teams, depending on the nature of the event.³³ Similarly, if an event is primarily spectator-based (eg, a football match in a stadium with assigned seats), considerations related

to site access and egress issues become highly relevant due to the not infrequent occurrence of riots and stampedes. Access and egress between the event and emergency services, as well as between the event and local hospitals or clinics, should also be a consideration.

Evont		
Boundaries	Venue/Site	Examples
Bounded	Venue Permanent	Stadium
	Venue Temporary	Stage Set Up for a
		Season
	Venue Mixed	National
		Exhibition
Unbounded	Single Site	Park
	Multiple Sites, Single Town/City	Community Days
	Multiple Sites, Shifting Footprint	Marathon
Compound	Multiple Sites, Multiple Towns/Cities. Event contains both sites directly involved in event and "nonevent spaces" between venues.	

Event Size—Early work by researchers in the field of MGH began with describing and classifying MGs according to the

Table 3. Classification of Mass Gatherings by Event Boundaries and Venue

Turris & 2014 Prehospital and Disaster Medicine

nmon Examples in Range	
Adventure obstacle race, triathlon, bike race, small venue music	
performance.	
Local music festival, charity fund-raiser sporting event, Ironman triathlon.	
Arts festival, regional fair, rodeo, community running event.	
Large marathon, out-door concert/music festival.	
Community celebration around national play-off game, outdoor music	
festival, national holiday.	
Parade, destination international music events.	
Fire-works display, protest march.	
Religious festivals and pilgrimages, international Games.	
International sport competitions, religious events (includes spectators,	
participants, and staff).	
Religious pilgrimage (ie, Hajj).	

 Table 4. Classification of Mass Gatherings According to Event Size
 Turris & 2014 Prehospital and Disaster Medicine

absolute number of attendees and/or participants. Malloy and colleagues cast a wide net and proposed that MGs be categorized according to a variety of sizes, including small, medium, large, major, super, extreme, and mega.60 This work contributes an understanding of the vast range of sizes, from local community events that occur on a small scale to international events that involve movement of large numbers of people across international boundaries. Based on existing classification schemes with regard to size, the following synthesis is presented (Table 4). The authors acknowledge that there are challenges related to labeling a given event as small, medium, and large, as such designations are difficult to interpret without information about the size of the host community. Instead, a range is provided. The size of a given MG has been used to quantify risk. In Western Australia, the public health authority uses the size of a

MG as one variable contributing to risk assessment score.⁶¹ this work links the absolute size of an event to risk assessment, which is an important consideration in relation to the permitting, planning, and execution of specific events.

Ratio of Event to Host Population Size — Recently, attention has shifted toward defining MGs as a preplanned event held at a

specific location for a defined period of time that has the potential to strain planning and response resources.⁵² This definition is appealing as it takes into account the baseline response capacity of the host community. Understanding the size of the event crowd relative to the size of the baseline host community population may contribute to understanding and quantifying risk or anticipating the impact of the MG. This could be expressed as a ratio of the event population relative to the host population (Table 5), or the event-to-host population (EHP) ratio. Essentially, this may be calculated as the number of people in the event community (N_{HC}).

Figure 1 draws on the proposed MG population model outlined in Part 1 of this series and presents a description and



Turris & 2014 Prehospital and Disaster Medicine			
Figure 1. Population Model for Mass-gathering Health.			

Ratio Range	Example	
<0.1	Crowd of 500 in community of .5,000; rodeo in small town.	
	Crowd of 250,000 in community of 3 million; fireworks in large city.	
0.11–0.50	30,000 attend mountain bike festival in a community of 70,000.	
	1,000,000 visitors attend football tournament in a city of 4,000,000.	
0.50–1.0 10,000 attend botanical show in community of 15,000 pec		
	24 million attend religious pilgrimage in country of 29 million.	
>1.0-2.0	18,000 attend a music festival in a community of 12,000 people.	
>2.0	36,500 attend concert in a community of 15,000.	

Turris & 2014 Prehospital and Disaster Medicine

Table 5. Examples of MGs Described by the EHP Ratio Abbreviations: EHP, event-to-host population; MG, massgathering.

diagrammatic representation of the relative size of events, according to the size of the population attending (Table 6). Operationally and practically speaking, arising from the EHP ratio are considerations about response capacity. In other words, if the absolute size of an event exceeds the size of the host community, health infrastructure may need to be augmented. Further work is needed to delineate how response capacity might be objectively described and determined.

Temporal Considerations—For this class of data, a myriad of time considerations apply. For example, how long is a givenevent (eg, hours, days, or weeks)? Does the event run for several days? Is the event taking place during the daytime, at night, or both? Outdoor seasonal events carry particular risks according to the weather conditions. Also, consider the differences between recurring (eg, annual) and one- off events. For example, recurring events may have the benefit of a legacy of experienced clinicians, volunteers, and participants.

Dynamics of an Event

Crowd Mood—Political rallies, funerals of public figures, and protests are examples of event classifications that influence the risk profile of a given MG. This is because crowd mood and behavior will be directly influenced by the tone of a given event, perhaps making the crowd more volatile. Increased stress for individual participants may play a role in crowd volatility; also, the risks associated with events that may be targets for deliberate acts of terror shape the relationship between the purpose of the event and risk to attendees.⁹ Crowd mobility and density are sometimes used as proxies for crowd mood or indicators of risk (qualitative). Objective measures of crowd mood are not yet well developed for use in a MG context.¹⁹

Crowd Mobility—Events have different "shapes" that influence how a crowd of participants and/or spectators may be distributed through the event, ultimately shaping crowd dynamics. Understanding the event "flow" as it relates to crowd mood and behavior may be valuable for both researchers and clinicians. For example, in relation to crowd mobility at a music festival where there is a highly-active stage front, crowd surfing is more likely to occur and must be planned for.

Crowd Density—Crowd density is another important feature of event description. Crowd density is defined as the ratio between the number of persons in the area (eg, 2.8 people/square meter is considered to be medium density; Table 7).

Steffen et al argued that crowd density was a more important consideration than the absolute number of people present. Measuring density at MGs is currently done in a nonstructured or anecdotal way. However, at a specific event, the environment can be divided into different regions and the density may be measured in each region (eg, entrance to an

event and stage fronts).³⁹ Similarly, Sun and Qin suggested dividing the environment into different regions to reflect the intensity and spaces of the crowd.⁶³

Density affects people in many ways. Berk argued that density affects visibility of those within the crowd, and dense

crowds will automatically undercut the effect of crowd activity.⁶⁴ For individuals in high-density crowds, it is difficult to see more than a few neighboring individuals. Polus claimed that there is no dispute that higher crowd densities lead to a greater interaction between individuals, and it is the impact of this interaction that is of interest, as is its

impact on audience members during an event and how it might impact behaviour.⁶⁵ A vivid example of the effect of crowd density on illness and injury rates may be found in the literature of stampedes that occur during religious gatherings^{23,66} and other types of events.⁶⁷⁻⁶⁹

Abbreviation	Term	Definition
Ν	Number of People	Integer representing the total population of interest.
НС	Host Community	The community (or communities) hosting the event.
N _{HC}	Number of People in the Host Community	Integer representing the total population of the host community or communities, on a nonevent day.
N _{Ev}	Event	Integer representing the number of people attending the event, including attendees, spectators, talent, and workforce.
N _{PP}	Patient Presentations	Number of patients presenting during the event; usually presenting to on-site health services, but may include direct presentations to community health services.
N _{HCEv}	Host Community and Event Population	Host community at baseline plus the number of people attending or participating in the event.

Primarily Spectator, Primarily Participant, Mixed Events—People at an event belong to one of three groups: spectators, participants, and event crew (eg, volunteers, security, vendors,

Turris & 2014 Prehospital and Disaster Medicine

Table 6. Abbreviations and Definition of Terms, Population Model for MGH Abbreviation: MGH, mass-gathering health.

Label	Metric
Extremely High Density	5 m ²
High Density	4 m ²
Medium Density	3 m ²
Low Density	2 m ²
Very Low Density	1 m ²
, , ,	

Turris & 2014 Prehospital and Disaster Medicine

Table 7. Crowd Density Measurement

Design of an Event

Special Hazards—The WHO adopts an "all hazards" approach to risk management and disaster planning, which may be applied to examining and understanding event design. A hazard (ie, "any phenomenon with the potential to cause disruption or damage to people or their environment") creates a degree of risk (ie, "the probability of a harmful

consequence").⁷⁰ Adopting an all-hazards approach involves planning and developing a plan to address the range of risks and possible emergencies that might occur during a given MG. In the context of health care in general, clinicians think about "modifiable" and "nonmodifiable" factors; however, in the context of MGs, the majority of hazards may be modifiable to some degree, through careful event design. Consider the following examples.

Infectious Disease Exposure—The degree of risk presented by ID exposure during a MG is an active area of inquiry for researchers. What can be accomplished to reduce risk for attendees is clear in the example of The Hajj, an annual religious festival attended by millions in Saudi Arabia (a review of the health risks posed is detailed in Ahmed

et al).⁷¹Based on the data from this annual event, the Saudi's have instituted a series of proactive measures to limit morbidity and mortality, thus the government supports 24 hospitals with a capacity of roughly 5,000 beds. Medical clinics (N 5136) in the vicinity of the Hajj offer free medical care to all pilgrims, and there are 18 stationary and 21 mobile health care teams at the event itself. Airports have clinical examination rooms and any pilgrim suspected

of having a communicable disease is taken by ambulance to a dedicated 200-bed hospital.¹⁵

Event Weather and Environmental Conditions—Depending on the season, temperature, and humidity, illness and

injuries at a given outdoor event may increase.³⁰ For example, as described by Bruce, heat-related casualties can run in

the thousands when there is no available protection for participants.⁷² In July of 2006, during the annual Nijmegen March in Holland, thousands of people walked 100 miles over four days, and a reported 2,725 heat-related casualties presented for treatment during a single day when the temperature rose to 428C. Similarly, the temperature and the humidity index may contribute to the illness rates for marathons, air shows, and other outdoor events held when temperatures are high, or even indoor events with inadequate air conditioning and/or ventilation. In addition to reports of temperature-related illness, there have been mass-casualty incidents at MGs related to weather

conditions. For example, a lightning strike caused fatalities during a festival,⁷³ a hailstorm triggered a human stampede,⁷⁴

and windy conditions caused the collapse of a large tent.⁷⁵ Although climactic conditions can be predicted, weather conditions on a particular day are variable and may change throughout the day. These examples point to the need to anticipate the unexpected by considering the range of expected weather conditions in a particular region rather than focusing on a single day or days.

Alcohol and Drug Use—The presence of drugs and alcohol, whether available through on-site vendors or brought in concealed by patrons, is an important consideration and can be difficult to capture in a meaningful way. For example, at an electronic dance music festival, alcohol may be sold on site; however, attendees who are underage and know that their bags will be searched upon entry to the event may choose to imbibe right before the event (eg, while waiting in the queue to enter), increasing their risk of alcohol toxicity⁵⁵

Mechanical or Other Hazards—Mechanical and other hazards are rarely reported on directly in event case reports. Instead, these types of hazards (eg, small planes at airshows,⁷⁶ amusement park rides, livestock at a rodeo, and fireworks at community celebrations⁷⁷⁻⁸¹) only appear in the literature when a mass-casualty event occurs. Collecting data on events with special hazards, prospectively, may assist researchers in measuring the degree of risk presented by similar events, ultimately informing event design.

Protective Factors—Illness and injury prevention is an emerging topic of discussion in the MGH literature. Several researchers have argued that prevention efforts should be a focus during event planning. For example, Hewitt and colleagues argued that many injuries that occurred during a rock concert were preventable if planners reduced the distribution of items that could serve as missiles, and suggested that limiting the size of liquid containers could be

considered as one means of reducing overall alcohol intake.⁴⁵

Intrinsic hazards, present by virtue of participation in particular types (eg, participation in a marathon or cycling event increases the risk of musculoskeletal injury), might be, to some degree, modifiable through careful event design. For example, organizers of the New York Marathon (USA) send out education materials about the prevention of injuries and illnesses both before and after the event (Personal Communication, Dr. Stuart Weiss, November 2012). The effect of this intervention has not yet been tested prospectively. Embedding illness and injury prevention in the event model will help researchers collect data and report on strategies being used to reduce the health care burden of MGs.

Health promotion vis a vis MGs has similarly been absent from the MGH literature until relatively recently. One example of an event-related, illness-prevention strategy is the Dance Safe initiative, designed to provide information

to attendees of electronic dance music events.⁸² At a more granular level, water stations, shelter, shade, access to sunscreen or earplugs, and available "chill out zones" are all examples of event design that have yet to be prospectively tested with regard to the effect on illness and injury rates.

Thinking more broadly about the positive health effects of events, Tewari and colleagues reported thought-provoking findings around the longitudinal health benefits of Hindu pilgrimage event. They found that participation in the pilgrimage increased the well-being of participants and argued that events should be reframed as not only presenting health risks, but also presenting opportunities for health benefits.⁴⁶

On-site Health Services—Community stakeholders have a vested interest in ensuring adequate planning to prevent or minimize the health impact of MGs in their communities. One of the ways to minimize the burden on local health care

services is to offer risk-relevant, on-site first aid and/or advanced health services throughout a special event. The MG literature is replete with case reports detailing event medical services. Researchers have been working on determining the "right" ratio of health care providers to attendees/participants for a given type of event, required equipment and supplies, effective communication plans, necessary transportation assets, and the appropriate infrastructure to support on-site health care services at different types of events.

Host Community Burden—In order to mitigate the health care burden MGs may place on baseline community health services, researchers would benefit from a common pool of data with which to measure community impact. Arbon et al noted that published MG literature to date consists predominantly of observational studies and cohort studies (>58%),

most commonly focused on operations and Emergency Medical Services on site at the event (>48%).³ Case reports and case series provide snapshots of specific events, commonly reporting patient presentations and transfers to hospital on the day(s) the event occurred. Unfortunately, these reports will underestimate the true impact of MGs on local community resources. Case reports rarely capture data regarding increased workload on health resources in the days or hours before and/or after the event. As argued by Lund and colleagues, a full-operational analysis would account for both the

consumptive and disruptive effects of MGs on the host community's health resources.²⁵ Well-executed medical and safety plans for events with appropriate, comprehensive risk assessments and stakeholder engagement have the best chance of ameliorating the potential negative impact of MGs on communities.

Overlay of the Population Model

This report has proposed a series of analytic lenses through which people might better understand and describe MGs and their health effects. This set of lenses is drawn from existing literature and forms the beginnings of an event model. In a companion paper, the present authors developed a population model, also central to further theory development. When the population model overlays the event model, this may focus the attention of researchers and clinicians on the links between the event characteristics and the event population. For example, the application of the event and population models may focus attention on the host-to-event population ratio in a rural setting, as in the first example (Table 1). The inverse high EHP ratio should prompt clinicians to coordinate with local health service infrastructure to temporarily increase capacity.

Moreover, as was the case with the population model, there is also an overlay of time for the event model. Ideally, the event model should be considered pre, during, and post event with data collection taking place during three phases in time.

Future Implications and Limitations

None of the event lenses described operate independently. For example, considering crowd behavior and density data together would support taking action to reduce risk for attendees, ideally before a negative health event takes place. Aly talks about "improving" and "dampening factors." He posited that as factors such as temperature and overall noise levels rise, so too does the probability of aggression, making a strong case for considering the interactions

between event factors.⁸⁸

The contribution of the work in this report is in helping researchers and clinicians articulate and understand risk related to specific events and may ultimately inform the execution of safer events. A more expanded view takes into account the variable scale of events and considers the respective health, security, and environmental impacts of MGs. Event models and classifications must have room for growth and adaptation as the science clarifies operational and academic validity of categories, and so on. As the science underpinning MGH continues to develop, it will move beyond identifying important parameters and existing conceptualizations toward the development of a robust event model.

Numerous definitions and classifications for MG events appear in the published literature. Ultimately, the purpose of event models will be to describe events in as much detail as possible in order that researchers and others can make comparisons between similar events and across similar event types. Also, researching the relationship between event type and risk will require consistency of event classifications and categories.

Conclusions

Because there will always be an interaction between event type and the people who attend or participate in the event, both event models and population models are required.

By moving towards a common understanding of population and event modeling as it pertains to MGs, the academic community will be better able to develop methodologies to describe, evaluate, establish evidence-based minimum standards, and study interventions aimed at medical services provision, health promotion, injury and illness prevention, and surveillance.

Clarity in definitions and descriptions of host community, event, and patient populations in the context of MGs will permit consistent description and reporting in the international literature. Importantly, it will also permit a better understanding of injury and illness presentations and clarify the health service impacts on the various populations of interest. Operational planning for emergency response, health promotion, injury and illness prevention, and surveillance will be measurable against more precisely-defined populations.

Research in Context

Literature Review

The writing team consisted of an international collaborative group with substantial experience with clinical, research, and policy development in the context of MGs. Review of the literature and addressing gaps in theory building identified in published reviews of the literature prompted the consensus process.

Interpretation

This manuscript puts forward theory-building concepts and models that may stimulate further discussion and consensus building amongst MG researchers in the international community.

Acknowledgements

The authors would like to acknowledge: Dr. Greg Anderson, PhD, Dean, Office of Applied Research, Justice Institute of BC, New Westminster, BC, Canada; Dr. Katherine Arbuthnott, WHO and Public Health England; and Dr. Mark Salter, Public Health England.

Authors' Contributions

Paul Arbon participated in the discussions at the Flinders University meeting and on team conference calls discussing the model and the publication; he also commented on the various drafts of the document. Ron Bowles contributed significantly to the theoretical modelling and, in conjunction with the principle author, expanded the paper into two parts in order to clarify the population of events vs the populations of people affected by mass gatherings. Elizabeth Ellison conducted the literature review for the two companion papers; she also commented on drafts of the document. Alison Hutton contributed to the initial draft of this paper and wrote the section on crowd dynamics; she was also part of the panel that gave feedback to the initial diagram concepts for this paper. Adam Lund cowrote the paper with the principle author; he also created the initial draft and the diagram concepts and was involved in synthesizing and editing all drafts of the manuscript. Consistent event description and modeling has been a focus of the University of British Columbia Mass-gathering Medicine Interest Group since 2009. Jamie Ranse contributed to the initial draft of this paper. Malinda Steenkamp participated in discussions at the Flinders University meeting and on team conference calls; he also contributed to the initial draft of this paper. Malinda Steenkamp participated in discussions at the Flinders University meeting and on team conference calls discussing the model and the publication; she also commented on the various drafts of the document. Sheila Turris is the principle author on this paper; she coordinated the team process and brought the paper to completion. She is also a member of the panel that created the original drafts of the figures and tables.

References

- 1. Arbon P. Introduction to the compendium on mass gatherings. *Prehosp Disaster Med*. 2010;25(6):1-2.
- 2. Arbon P. The development of conceptual models for mass-gathering health. *Prehosp Disaster Med*. 2004;19(3):208-212.
- 3. Arbon P, Cusack L, Verdonk N. Mass gathering public health and emergency medicine literature review: levels of evidence. *Austalasian J of Paramed*. 2013;10(1): 1-5.
- 4. Memish ZA. Jeddah declaration on mass gatherings health. Lancet. 2011;11(5): 342-343.
- 5. Ranse J, Hutton A. Minimum data set for mass gathering health research and evaluation: a discussion paper. *Prehosp Disaster Med*. 2012;27(6):1-8.
- 6. Turris SA, Lund A. Minimum data set for mass-gatherings health research and evaluation: a response. *Prehosp Disaster Med*. 2013;28(2):191-193.
- 7. Lund A, Turris SA, Bowles R, et al. Mass-gathering health research foundational theory: part 1 population models for mass gatherings. *Prehosp Disaster Med*. In press.
- 8. Alquthami AH, Pines JM. A systematic review of noncommunicable health issues in mass gatherings. *Prehosp Disaster Med*. 2014;29(2):1-9.
- 9. Steffen R, Bouchama A, Johansson A, et al. Non-communicable health risks during mass gatherings. *Lancet*. 2012;12(2):142-149.
- 10. Abubakar I, Gautret P, Brumnette GW, et al. Global perspectives for prevention of infectious diseases associated with mass gatherings. *Lancet*. 2012;12(2):66-74.
- 11. Botelho-Nevers E, Gautret P. Outbreaks associated to large open air festivals, including music festivals, 1980-2012. *Eurosurveill*. 2013;18(11):1-10.
- 12. Centers for Disease Control and Prevention Web site. http://wwwnc.cdc.gov/travel/ yellowbook/2014/chapter-8-advising-travelers-with-specific-needs/travel-to-mass- gatherings. Accessed November 13, 2013.
- 13. Chowell G, Nishiura H, Viboud C. Modeling rapidly disseminating infectious disease during mass gatherings. BMC Med. 2012;10:159-169.
- 14. Khan K, McNabb SJN, Memish ZA, et al. Infectious disease surveillance and modeling across geographic frontiers and scientific specialties. *Lancet*. 2012;12(3):222-230.
- 15. Memish ZA. The Hajj: communicable and non-communicable health hazards and current guidance for pilgrims. *Euro Surveill*. 2010;15(39):196.
- 16. Shafi S, Booy R, Haworth E, Rashid H, Memish ZA. Hajj: health lessons for mass gatherings. *J Infection Public Health*. 2008;1(1):27-32.
- 17. Tam JS, Barbeschi M, Shapovalova N, Briand S, Memish ZA, Kieny MP. Research agenda for mass gathering: a call to action. *Lancet*. 2012;12(2):231-239.
- 18. Johansson A, Batty M, Hayashi K, Al Bar O, Marcozzi D, Memish ZA. Crowd and environmental management during mass gatherings. *Lancet*. 2012;12(2):150-156.
- 19. Memish ZA. Emergency of medicine for mass gatherings: lessons from the Hajj. Lancet. 2012;12(1):56-65.
- 20. Soomaroo L, Murray V. Disasters at mass gatherings: lessons from history. PLOS Currents. 2012;4:1-12.
- 21. Turris SA, Lund A, Bowles R. Mass casualty incidents at mass gatherings a review of the literature. *Disaster Med Pub Health Preparedness*. 2014;7(1):1-7.
- 22. Begum AA. Unnatural deaths during Zakat distribution. *Bangladesh Med Res Conc Bull*. 1993;19(3):99-102.
- 23. Hsieh YH, Ngai KM, Burkle FM, Hsu EB. Epidemiological characteristics of human stampedes. *Dis Med Public Health Prep*. 2009;3(4):217-223.
- 24. Jangi S. Under the medical tent at the Boston Marathon. NEJM. 2013;368(21): 1953-1955.
- 25. Lund A, Turris SA, Bowles R. Conceptualizing the impact of special events on community health service levels: an operational analysis. *Prehosp Disaster Med*. 2014;29(5):525-532.
- 26. Mahoney EJ, Harrington DT, Biffl WL, Metzger J, Oka T, Cioffi WG. Lessons learned from a nightclub fire: institutional disaster preparedness. *J Trauma*. 2005;58(3):487-491.
- 27. Hnatow DA, Gordon DJ. Medical planning for mass gatherings: a retrospective review of the San Antonio Papal Mass. *Prehosp Disaster Med*. 1991;6(4):443-450.
- 28. Kenar L, Pharm TK. Medical preparedness against chemical and biological incidents for the NATO Summit in Istanbul and lessons learned. *Prehosp Disaster Med.* 2006;21(4):268-271.

- 29. Leonard RB, Winslow JE, Bozeman WP. Planning medical care for high-risk mass gatherings. *Internet J of Rescue* & *Disaster Med*. 2007;6(1):1-15.
- 30. Milsten AM, Maguire BJ, Bissell RA, Seaman KG. Mass-gathering medical care: review of the literature. *Prehosp Disaster Med*. 2002;17(3):151-162.
- 31. Nguyen RB, Milsten AM, Cushman JT. Injury patterns and levels of care at a marathon. *Prehosp Disaster Med*. 2008;23(6):19-25.
- 32. Rose WD, Larid SL, Prescott JE. Emergency Medical Services for collegiate football games: a six and one-half year review. *Prehosp Disaster Med*. 1992;7(2):157-159.
- 33. Turris SA, Lund A, Gutman SJ, Mui J, Wang P, Lewis K. An organized medical response for the Vancouver International Marathon (2006-2011): when the rubber hits the road. *Cur Sports Med Reports*. 2014;13(3):147-154.
- 34. Yazawa K, Kamijo Y, Sakai R, Ohashi M, Owa M. Medical care for a mass gathering: the Suwa Onbashira Festival. *Prehosp Disaster Med*. 2007;22(5):431-435.
- 35. Zeitz KM, Schneider D, Jarrett D. Mass gathering events: a retrospective analysis of patient presentations over seven years. *Prehosp Disaster Med*. 2002;17(3):147-150.
- 36. Feldman MJ, Lukins JL, Verbeek RP, MacDonald RD, Burgess RJ, Schwartz B. Half a million strong: the emergency medical services response to a single-day, mass- gathering event. *Prehosp Disaster Med*. 2004;19(4):287-296.
- 37. Zielioski A, Pawlak BJ. Toolbox for implementation of surveillance at mass gatherings. WP 4: surveillance during mass gatherings. http://www.rki.de/EN/ Content/Prevention/React/Work/wp4/WP_4_ToolBox.pdf? blob5publicationFile. Accessed May 1, 2014.
- 38. Arbon P. Planning medical coverage for mass gatherings in Australia: what we currently know. *J Emerg Nurs*. 2005;31(4):346-350.
- 39. Hutton A, Brown S, Verdonk N. Exploring the role of culture as a psychosocial motivator at mass gathering events. *Prehosp Disaster Med*. 2009;28(5):292-297.
- 40. Morimura N, Katsumi A, Koldo Y. Analysis of patient load data from FIFA World Cup Korea/Japan. *Prehosp Disaster Med*. 2004;19(3):278-284.
- 41. Pinkert M, Bloch Y, Schwartz D, et al. Leadership as a component of crowd control in a hospital dealing with a mass-casualty incident: lessons learned from the October 2000 riots in Nazareth. *Prehosp Disaster Med*. 2007;22(6):522-526.
- 42. Salhanick SD, Sheahan W, Bazarian JJ. Use and analysis of field triage criteria for mass gatherings. *Prehosp Disaster Med*. 2003;18(4):347-352.
- 43. Emergency Management Australia. Safe and healthy mass gatherings: a health, medical and safety planning manual for public events. http://www.nationalcapital.gov. au/downloads/events_and_venues/assessment_criteria/ESA_Mass_gatherings.pdf. Accessed May 1, 2014.
- 44. Gutman SJ, Lund A, Turris SA. Mass medical support for the 2009 World Police and Fire Games: a descriptive analysis of a large scale participation event and its impact. *Prehosp Disaster Med*. 2011;26(1):33-39.
- 45. Hewitt S, Jarrett L, Winter B. Emergency medicine at a large rock festival. J Accid Emerg Med. 1996;13(1):26-27.
- 46. Tewari S, Khan S, Hopkins N, Srinivasan N, Reicher S. Participation in mass gatherings can benefit well-being: longitudinal and control data from a North Indian Hindu pilgrimage event. *PLOS One*. 2012;7(10):1-5.
- 47. Zeitz KA, Zeitz C, Arbon P, Cheney F, Johnston R, Hennekam J. Practical solutions for injury surveillance at mass gatherings. *Prehosp Disaster Med*. 2008;23(1):76-81.
- 48. Bledsoe B, Songer P, Buchanan K, Westin J, Hodnick R, Gorosh L. Burning man 2011: mass gathering medical care in an austere environment. *Prehosp Disaster Med*. 2012;16(4):469-476.
- 49. Ma O, Pirrallo R, Rubin J. Survey of medical services at major league baseball stadiums. *Prehosp Disaster Med*. 1995;10(4):268-271.
- 50. Milsten AM, Seaman KG, Lui P, Bissell RA, Maguire BJ. Variables influencing medical usage rates, injury patterns, and levels of care for mass gatherings. *Prehosp Disaster Med.* 2003;18(4):334-346.
- 51. Arbon PA. Mass gathering medicine: a review of the evidence and future directions for research. *Prehosp Disaster Med*. 2007;22(2):131-135.
- 52. Turris SA, Lund A. Triage during mass gatherings. Prehosp Disaster Med. 2012;27(6):531-535.

- 53. Erickson TB, Koenigsberg M, Bunney EB, et al. Prehospital severity scoring at major rock concert events. *Prehosp Disaster Med*. 1997;12(3):195-199.
- 54. Carmont MR, Sedgwick DM. The impact of an extreme sports event on a district general hospital. *Scot Med J*. 2005;50(3):106-108.
- 55. Lund A, Turris SA. Electronic dance music events (EDME) risks, patient presentations and public health care costs. Submitted to: *Canadian J of Emergency Med*. (in review).
- 56. Giulianotti R. Football and the politics of carnival: an ethnographic study of Scottish fans in Sweden. *Int Rev Socio Sport*. 2013;48(5):191-220.
- 57. Marsh P, Fox K, Carnibella G, McCann J, Marsh J. Football violence and hooliganism in Europe. 1996. http://www.sirc.org/publik/fvexec.html. Accessed May 14, 2013.
- 58. Peiterson B. Supporter culture in Denmark: the legacy of the 'world's best supporters'. *Soccer and Society*. 2009;10(3-4):374-385.
- 59. Stott C, Adang O. Understanding and managing risk: policing football matches. *Bavnekbanke Press*. 1995;995:7-14.
- 60. Malloy MS, Sherif Z, Natin S, McDonnell J. "Management of mass gatherings." In: Koenig KL, Shultz CH. *Disaster Medicine: Comprehensive Principles and Practices*. Cambridge, United Kingdom: Cambridge University Press, 2010.
- 61. Public Health Western Australia. http://www.public.health.wa.gov.au/cproot/2542/ 2/Medical%20Risk%20Classification.pdf. Accessed February 25, 2014.
- 62. World Health Organization. 2008. Communicable disease alert and response: key considerations. http://www.who.int/csr/Mass_gatherings2.pdf. Accessed February 26, 2014.
- 63. Sun L, Li X, Wenhu Q. Simulating realistic crowd based on agent trajectories. *Computer Animation Virtual Worlds*. 2013;24(3-4):165-172.
- 64. Berk RA. A gaming approach to crowd behavior. American Sociol Rev. 1974;39:355-373.
- 65. Polus A, Schofer JL, Ushpiz A. Pedestrian flow and level of service. *J Transp Eng.* 1983;109:46-56.
- 66. Ngai KM, Burkle FM, Hsu A, Hsu EB. Human stampedes: a systematic review of historical and peer-reviewed sources. *Dis Med Pub Health Prep*. 2009;3(4):191-195.
- 67. Ackermann O, Lahm A, Pfohl M, et al. Patient care at the 2010 love parade in Duisburg, Germany. *Deutsches Arzteblatt International*. 2011;108(28-29):483.
- 68. Bemelman M, Leenen L. Mass casualty event during a musical parade: lessons learned. *European J Trauma Emerg Surg*. 2008;5:451-456.
- 69. Helbing D, Johansson A, Al-Abideen HZ. The dynamics of crowd disasters: an empirical study. *Phys Rev*. 2007;75(4):1-7.
- 70. World Health Organization. Risk reduction and emergency preparedness. 2007. http://www.who.int/hac/techguidance/preparedness/emergency_preparedness- s_eng.pdf. Accessed April 11, 2014.
- 71. Ahmed QA, Arabi YM, Memish ZA. Health risks at the Hajj. Lancet. 2006;367(9515):1008-1015.
- 72. Bruce ASW, Tunstall C, Boutler MJ, Coneybeare A. A treatment algorithm for mass heat casualties. *JF Army Med Corps*. 2008;154(1):19-20.
- 73. 17 injured in lightning strike at Toronto food festival. http://www.cbc.ca/ news/canada/toronto/story/2012/07/15/toronto-whitby-lightning-strike.html. Accessed February 25, 2013.
- 74. 93 die in Nepal stadium stampede: soccer fans rush to locked exits in sudden hailstorm. http://news.bbc.co.uk/2/hi/uk_news/768380.stm. Accessed April 2, 2013.
- 75. Deal KE, Synovitz CK, Goodloe JM, King B, Stewart CE. Tulsa, Oklahoma Oktoberfest ten collapse report. *Emerg Med Int*. 2012:1-9.
- 76. Howell W. Airshow calamity tests Reno's hospital preparedness. *Hosp Health Network J.* 2011;85(12):16-17.
- 77. Spanish church fireworks display explodes. CBC News. August 14, 2012. http:// www.cbc.ca/news/world/story/2012/08/14/accidental-fireworks.html. Accessed May 14, 2013.
- 78. Fireworks explode in crowd in China. Telegraph. October 14, 2012. http:// www.telegraph.co.uk/news/worldnews/asia/china/9607552/Fireworks- explode-in- crowd-in-China.html. Accessed February 27, 2013.

- 79. Several injured during Liberty Park fireworks show. July 25, 2010. http:// www.abc4.com/news/local/story/Several-injured-during-Liberty- Park-fireworks- show/mxloGVnciUewhxWMBLG8 g.cspx. Accessed February 27, 2013.
- 80. Fireworks explosion kills 21. http://ntdtv.org/en/news/china/ 2010-03-01/ 040965832541.html. Accessed February 27, 2013.
- Suphan Buri fireworks display affects 500. *The Nation*; January 26, 2012. http://www.nationmultimedia.com/national/Suphan-buri-fireworks-disaster-affects- 500-30174526.html. Accessed February 27, 2013.
- 82. Dance safe. http://www.dancesafe.org/. Accessed April 18, 2014.
- 83. Kerr GW. Emergency medical planning at soccer matches. Int J Sports Med. 2003;4(1):1-5.
- 84. Engledow S, Yancey A. EMS to the extreme. Firechief. 2012;30-36.
- 85. Rubin AL. Safety, security, and preparing for disaster at sporting events. *Curr Sports Med Rep*. 2004;3(3):141-145.
- 86. Arbon P, Bridgewater FHG, Smith C. A predictive model for patient presentation and transport rates. *Prehosp Disaster Med*. 2001;16(3):150-158.
- 87. Zeitz KM, Zeitz CJ, Arbon P. Forecasting medical work at mass-gathering events: predictive model versus retrospective review. *Prehosp Disaster Med.* 2005;20(3):164-168.
- 88. Aly M. Dynamics of riots. Resilence. 2013:32-34.